

Representations of the caregiver–child relationship and of the self, and emotion regulation in the narratives of young children whose mothers have borderline personality disorder

JENNY MACFIE AND SCOTT A. SWAN

University of Tennessee at Knoxville

Abstract

Borderline personality disorder (BPD) represents a severe distortion in the development of attachment, self, and emotion regulation. Study of children at high risk of developing BPD may inform precursors to BPD. In a low socioeconomic status sample of 30 children aged 4–7 whose mothers have BPD and 30 normative comparisons, representations of the caregiver–child relationship and of the self, and emotion regulation were assessed with a story-stem completion measure. In contrast to comparisons and controlling for major depressive disorder, children whose mothers have BPD told stories with the following: (a) more parent–child role reversal, more fear of abandonment, and more negative mother–child and father–child relationship expectations; (b) more incongruent and shameful representations of the self; and (c) poorer emotion regulation indicated by more confusion of boundaries between fantasy and reality and between self and fantasy, more fantasy proneness, less narrative coherence, and marginally more intrusion of traumatic themes. In the sample as a whole, (a) a maladaptive caregiver–child relationship composite was associated with maternal identity disturbance and self-harm; (b) a maladaptive self-composite was associated with maternal self-harm; and (c) a maladaptive emotion regulation composite was associated with maternal identity disturbance, negative relationships, and self-harm. Results are discussed in terms of putative precursors to BPD and preventive interventions.

If a mother experiences extreme emotional turbulence, intense unstable relationships, and confusion in her sense of self, and if she engages in impulsive behavior (e.g., with alcohol or drugs), suicidal gestures, self-mutilation, and inappropriate outbursts of anger, how may these affect her child's development? Because borderline personality disorder (BPD) has been character-

ized as a disorder of attachment, self-development, and emotion regulation, domains conceptually similar to the developmental tasks of early childhood (Sroufe, Egeland, Carlson, & Collins, 2005), a mother who has BPD may present a uniquely challenging context for her child's development (Macfie, 2009). It is unlikely that BPD develops spontaneously in early adulthood so we need to understand developmental precursors (Crick, Murray-Close, & Woods, 2005). Precursors may include mental representations carried forward from early childhood (Carlson, Sroufe, & Egeland, 2004), which increase the possibility of the development of BPD in early adulthood. The current study examines representations of the caregiver–child relationship and of the self,

This research was supported by Grant MH077841 from the National Institute of Mental Health (to J.M.). We are grateful to the participating families for their contributions of time and effort.

Address correspondence and reprint requests to: Jenny Macfie, 301E Austin Peay, University of Tennessee, Knoxville, TN 37996-0900; E-mail: macfie@utk.edu.

and emotion regulation in the stories of children age 4–7 whose mothers have BPD.

Developmental Psychopathology

From a developmental psychopathology perspective (Cicchetti, 1984; Sroufe & Rutter, 1984), an understanding of pathways toward and away from disorder may inform adaptive development and preventive interventions (Cicchetti & Hinshaw, 2002; Cicchetti & Toth, 1992, 2006). To examine these pathways we need to study children at high risk of developing the disorder. Offspring of mothers with BPD comprise one of several high-risk groups for developing BPD (Lenzenweger & Cicchetti, 2005). Indeed, children are at high risk of developing their mothers' disorder (Downey & Coyne, 1990; Mednick & McNeil, 1968) because of a transaction between environmental and biological factors (Sameroff & Chandler, 1975). There is a very large body of research on children of depressed mothers that has informed successful preventive interventions (Cicchetti, Rogosch, & Toth, 2000; Cicchetti, Toth, & Rogosch, 1999; Toth, Rogosch, Manly, & Cicchetti, 2006). There are, by contrast, only six studies of children whose mothers who have BPD, reviewed below (Barnow, Spitzer, Grabe, Kessler, & Freyberger, 2006; Crandell, Patrick, & Hobson, 2003; Feldman et al., 1995; Hobson, Patrick, Crandell, Garcia-Perez, & Lee, 2005; Newman, Stevenson, Bergman, & Boyce, 2007; Weiss et al., 1996).

BPD

A BPD diagnosis cannot be made officially until early adulthood (American Psychiatric Association, 1994), but it may be possible in adolescence (Ludolph et al., 1990). BPD requires five or more of symptoms grouped here under the domain of dysfunction. First, BPD has been characterized as a disorder of *attachment* (Fonagy, Target, & Gergely, 2000; Gunderson, 1996; Liotti & Pasquini, 2000), with symptoms including fear of abandonment, volatile relationships, and inappropriate angry outbursts. Second, BPD has been characterized as a disorder of *self-development* (Westen & Cohen, 1993), with symptoms including an unstable sense of identity, feelings of emptiness, and brief paranoid or dissociative states.

Third, BPD has been characterized as a disorder of *emotion regulation* (Posner et al., 2003), with symptoms including self-damaging impulsivity, suicidal behaviors and/or self-injury, emotional reactivity, and inappropriate displays of anger. A diagnosis of BPD thus includes symptoms that reflect relationship and identity disturbance that may be more experience based and symptoms that reflect poor emotion regulation that may be more biologically based (Lenzenweger & Cicchetti, 2005).

In a large nationally representative study, 5.9% of the population were found to have BPD (Grant et al., 2008), a prevalence that is twice as high as formerly reported (American Psychiatric Association, 1994; Swartz, Blazer, George, & Winfield, 1990). Moreover, functioning in these individuals is as low as it is for those with schizophrenia (Gunderson, Carpenter, & Strauss, 1975). Furthermore, although equally prevalent in men and women (not more common in women as previously thought), BPD is associated with more severe mental and physical disability in women (Grant et al., 2008). Completed suicide occurs in 8–10% of individuals with BPD (American Psychiatric Association, 1994; Paris, 1993; Stone, 1990).

BPD is thought to develop from an interaction between an emotionally vulnerable child and an emotionally unsupportive environment (Heard & Linehan, 1993). Emotional vulnerability may lie in BPD's large hereditary component (Torgersen et al., 2000), perhaps in temperamental variables such as impulsivity and affective instability (Siever & Davis, 1991). An emotionally unsupportive environment may include child maltreatment and separation from or loss of caregivers (Herman, Perry, & Kolk, 1989; Laporte & Guttman, 1996; Weaver & Clum, 1993; Zanarini, 2000) and invalidating reactions to the child's needs and feelings (Heard & Linehan, 1993).

Success or failure at negotiating early developmental issues of *attachment* in the first year, *self-development* (autonomy) in the toddler period, and *self-regulation* (of emotion and behavior) in the preschool period (Sroufe & Rutter, 1984) may make the development of BPD more likely. Success or failure at each developmental issue depends in part upon success or failure at the previous one, so initial failure may have a cascading effect (Cicchetti, 1984).

Prior Research on BPD Offspring

Mothers with BPD are more intrusively insensitive when their infants are 2 months, and their infants look more dazed and more depressed than do normative comparisons (Crandell et al., 2003). At 13 months, these mothers with BPD are still more intrusively insensitive and infants are still more depressed, plus 80% are classified as disorganized in their attachment (Hobson et al., 2005). In similar findings, mothers with BPD of infants aged 3–36 months are less sensitive, and their infants are less eager to interact than are normative comparisons (Newman et al., 2007).

For children sampled between the preschool period and adolescence whose mothers had BPD, families are more unstable and more unhappy (Feldman et al., 1995), and children have more impulse control disorders compared with a normative sample (Weiss et al., 1996). Furthermore, for children age 11–18 whose mothers have BPD, mothers are more overprotective and the children demonstrate more emotional and behavioral problems than do normative and clinical comparisons (Barnow et al., 2006).

Internal Working Models

Internal working models are theorized to develop within the attachment relationship between infant and caregiver such that representations of others and of the self are used to guide expectations and behavior in future relationships (Bowlby, 1969/1982; Bretherton & Munholland, 2008). If the caregiver is sensitive and responsive, attachment is classified as secure and the infant is thought to develop an internal working model of others as trustworthy and of the self as worthy of care; if the caregiver is unavailable or inconsistently available, attachment is classified as insecure, and the infant is thought to develop an internal working model of others as rejecting or ambivalent and of the self as not worthy of care (Ainsworth, Blehar, Waters, & Wall, 1978). If the caregiver is frightening, the child is unable to develop an organized attachment and is classified as disorganized (Hesse & Main, 2006; Main & Solomon, 1990). A child disorganized in infancy may develop an internal working model of others as frightening, and of the self as bad and incongruent.

For early experience to affect development beyond heritable variables and the current environment, experience has to be internalized and carried forward (Carlson et al., 2004). Internal working models may provide the process. Although internal working models cannot be viewed directly, following the development of language they may be assessed at the level of representation using semiprojective measures such as the completion of the beginnings of stories (Bretherton & Munholland, 2008; Bretherton, Ridgeway, & Cassidy, 1990; Main, Kaplan, & Cassidy, 1985). Indeed, representations assessed in each developmental period between preschool and adolescence were significantly correlated with each other and with previous, current, and subsequent teacher-rated behavior (Carlson et al., 2004).

Internal working models carried forward from early childhood may contribute to the development of BPD. We know that representations of adults with BPD are more malevolent and less empathic than are those of comparisons (Nigg, Lohr, Westen, Gold, & Silk, 1992). Did these representations originate in early childhood? We also know that infant–mother attachment tends to be disorganized when the mother has BPD (Hobson et al., 2005), and that disorganized attachment develops if the caregiver is frightening (Hesse & Main, 2006; Main & Solomon, 1990). The infant is unable to find security in the relationship, and is caught between approach and avoidance and disorganizes under stress. To gain some feeling of security in the relationship, this child may develop a parent–child role reversal that has thus far been observed both in the toddler period (Macfie, Fitzpatrick, Rivas, & Cox, 2008) and at age 6 (Main et al., 1985). Moreover, role reversal in the toddler period in turn predicts behavior problems in kindergarten (Macfie, Houts, McElwain, & Cox, 2005) and is transmitted intergenerationally (Macfie, McElwain, Houts, & Cox, 2005).

Narrative Story-Stem Completions

A story-stem completion measure (Scarlett & Wolf, 1979) has been utilized to assess narrative representations in children age 3–7. Children complete the beginnings of stories presented to them by an examiner about conflictual or other

emotionally charged themes in family life. The stories are presented in a dramatic fashion with family figures and props, and because this is consistent with developmentally appropriate play, children enjoy completing them. The resulting narratives may be coded for representations of the caregiver–child relationship and of the self-thought to reflect internal working models (Bretherton, Ridgeway, et al., 1990; Macfie et al., 1999; Toth, Cicchetti, Macfie, Maughan, & VanMeeuwen, 2000).

In addition to assessing *representations*, creating and transmitting a narrative in the face of challenging emotional themes may reflect important aspects of *emotion regulation* (Oppenheim, 2006; Oppenheim & Waters, 1995). Emotion regulation allows for flexible and adaptive ways to meet challenges that individuals with BPD often lack (Putnam & Silk, 2005), and that contributes to resilience in at-risk children (Curtis & Cicchetti, 2007). Emotion regulation is thought to originate in the attachment relationship, specifically in how well the caregiver communicates with the child (Bowlby, 1988; Main et al., 1985). Thus, the attachment relationship is thought to be the source of not only internal working models but also of emotion regulation, both of which may be assessed in children's narratives.

One component of emotion regulation in the context of completing story stems is the ability to maintain boundaries around the narrative frame, boundaries that are newly developed in the preschool period and may be temporarily lost if the child experiences strong feelings such as excitement and fear (Scarlett & Wolf, 1979). These include the ability to distinguish between what is real and what is fantasy, and the ability to stay outside the story rather than becoming an actor in it. Another is the child's ability to create coherent narratives that address the issues presented (Oppenheim, Nir, Warren, & Emde, 1997; Oppenheim & Waters, 1995), and that do not veer off into irrelevant fantasy or disorganize into chaotic traumatic themes.

Validity of the Narrative Story-Stem Measure

What do narrative representations from story-stem completions reflect? There is considerable evidence that narratives reflect internal working

models of what actually happens in the child's life. In the review that follows, children are sampled in the preschool period unless otherwise stated.

Representations of caregivers

In normative samples, narrative representations of caregivers reflect attachment classifications assessed in infancy, suggesting continuity of internal working models (Bretherton, Ridgeway et al., 1990; Main et al., 1985; Solomon, George, & DeJong, 1995). Narrative representations of caregivers also correspond to maternal distress (Oppenheim, Emde, & Warren, 1997) and to children's independence, self-esteem (Oppenheim, Nir et al., 1997), anxiety (Warren, Emde, & Sroufe, 2000), prosocial behavior, and conduct problems (Stadelmann, Perren, Wyl, & Klitzing, 2007).

In at-risk samples, narrative representations of caregivers distinguish between maltreated and nonmaltreated children on responses to child distress (Macfie et al., 1999). Narrative representations of caregivers are also associated with child depression (Belden, Sullivan, & Luby, 2007), and become more positive and less negative following child–parent psychotherapy (Toth, Maughan, Manly, Spagnola, & Cicchetti, 2002). Moreover, exposure to domestic violence is associated with aggression and avoidance in narratives (Grych, Wachs-muth-Schlaefler, & Klockow, 2002), which varies according to the severity of mothers' post traumatic stress symptoms (Schechter et al., 2007). Furthermore, maternal depression in the toddler period predicts an increase of negative and a decrease of positive representations of parents between age 3 and age 4, controlling for recent and concurrent depressive symptoms (Toth, Rogosch, Sturge-Apple, & Cicchetti, 2009).

Representations of self

In normative samples, narrative representations of children reflect the child's hostility, hyperactivity (Zahn-Waxler, Schmitz, Fulker, Robinson, & Emde, 1996), and externalizing problems (Warren, Oppenheim, & Emde, 1996). In at-risk samples the experience of having

been maltreated is associated with negative child representations (Toth, Cicchetti, Macfie, Maughan, et al., 2000), more conflictual themes, fewer moral-affiliative themes (Toth, Cicchetti, Macfie, Maughan, 2000), fewer child responses to the distress of other children, and more responses to the distress of parents in a role reversal (Macfie et al., 1999). Furthermore, antisocial boys demonstrate more aggression and less understanding of the feelings or motives of others (Hill, Fonagy, Lancaster, & Broyden, 2007), whereas social competence in school is associated with prosocial themes (vonKlitzing, Stadelmann, & Perren, 2007). Moreover, in middle childhood disorganization in narratives is associated with externalizing symptoms (Green, Stanley, & Peters, 2007), and a composite of maltreated children's representations and behavior reflects more putative precursors to BPD than for comparisons (Rogosch & Cicchetti, 2005). Finally, and perhaps most importantly, maltreated children's representations of conflictual themes partially mediate the relationship between maltreatment and externalizing symptoms (Toth, Cicchetti, Macfie, Rogosch, et al., 2000), supporting a theory of internal working models as the process.

Emotion regulation in narratives

In normative samples, narrative coherence is predicted both by secure infant–mother attachment (Bretherton, Ridgeway, et al., 1990), and by emotional coherence in the mother–child relationship, and it is negatively correlated with externalizing symptoms (Oppenheim, Nir, et al., 1997). In addition, children who were classified as disorganized in infancy display deficits in emotion regulation with the intrusion of traumatic or bizarre themes in their narratives (Cassidy, 1988). Narrative coherence also has biological correlates. Children with lower vagal tone, when responding to an emotionally laden story stem versus a neutral story stem, tell more coherent stories than do those with increased vagal tone; and stories involving separation/reunion lead to cardiac responses that reflect increased emotional load (Bar-Haim, Fox, VanMeenen, & Marshall, 2004).

In at-risk samples, maltreated children confuse self and fantasy, confuse reality and fan-

tasy, display fantasy proneness, and the intrusion of traumatic material. All are associated with dissociation, an ineffective means of emotion regulation (Macfie, Cicchetti, & Toth, 2001). Furthermore, children age 4–7 exposed to domestic violence display lower narrative coherence (Schechter et al., 2007). Moreover, maltreated children in middle childhood demonstrate low narrative coherence in a maladaptive composite that is associated with emotional dysregulation and peer rejection (Shields, Ryan, & Cicchetti, 2001).

The Current Study

What might we expect in the narratives of children whose mothers have BPD under the stress of completing stories with emotionally charged family themes? Because most of these children have a history of disorganized attachment (Hobson et al., 2005), representations of the caregiver–child relationship may include role reversal (Macfie et al., 2008; Main et al., 1985), fear of abandonment, and negative parent–child relationship expectations. A composite of these maladaptive caregiver–child relationship representations may be associated with those maternal features of BPD that affect relationships: identity disturbance, negative relationships, and self-harm.

Corresponding self-representations may be more incongruent in the context of the mother's own inconsistent behavior (Bezirgianian, Cohen, & Brook, 1993) and more negative and shameful in the context of a mother who may have grown up feeling that her own emotional reactions were invalid (Heard & Linehan, 1993). A composite of maladaptive self-representations may be associated with those maternal features of BPD that relate to hurting oneself: self-harm.

Emotion regulation develops ideally in the context of a secure attachment (Bowlby, 1988; Main et al., 1985). Because offspring of women with BPD are likely to have been disorganized in their attachment, and because mothers with BPD themselves display severe emotional dysregulation, their children may display emotion dysregulation also. Under stress they may not be able to maintain boundaries around the narrative frame, confusing both self and fantasy and reality and fantasy (Scarlett & Wolf, 1979). They may also not be able to tell a coherent story, and veer off

instead into unrelated fantasy or chaotic traumatic themes. A composite of maladaptive emotion regulation may be associated with those features of maternal BPD that reflect poor emotion regulation: affective instability, identity disturbance, negative relationships, and self-harm.

In the current study, research on narrative representations of the caregiver–child relationship and of the self, together with emotion regulation, was extended to children age 4–7 whose mothers have BPD. Because of widespread comorbidity of BPD with major depressive disorder (MDD; Zanarini et al., 1998), and because MDD exerts such a large negative effect on child development (Cicchetti & Toth, 1998; Downey & Coyne, 1990), the effect of MDD was assessed as a covariate. We hypothesized that BPD offspring age 4–7 versus normative comparisons would (a) portray more maladaptive caregiver–child relationships in terms of more role reversal, more fear of abandonment, more negative mother–child and father–child relationship expectations; (b) portray more maladaptive self-representations in terms of more negative, more incongruent, and more shameful representations of children; and (c) demonstrate poorer emotion regulation in terms of more reality/fantasy confusion, more self/fantasy boundary confusion, more fantasy proneness, more intrusion of traumatic material and less narrative coherence.

In addition to assessing BPD categorically, BPD features were derived from maternal self-report and their association with children's narratives was assessed. It was hypothesized that in the sample as a whole, (d) a maladaptive child–caregiver relationship composite would correlate significantly with maternal identity disturbance, negative relationships, and self-harm; (e) a maladaptive self-composite would correlate significantly with maternal self-harm; and (f) a maladaptive emotion regulation composite would correlate significantly with maternal affective instability, identity disturbance, negative relationships, and self-harm.

Method

Participants

A low socioeconomic status (SES) sample consisted of a total of 60 children: 30 children whose

mothers had BPD, and 30 children whose mothers did not. There were 15 boys and 15 girls in each group, and 12% of the sample had a minority ethnic background (8.5% Hispanic, 3.5% African American). Their average age was 5 years, 4 months ($SD = 10$ months, range = 4 years, 0 months to 6 years, 11 months). The two groups were matched on child age and child verbal ability. Groups were also matched on demographic variables: family income, presence of partners, number of adults and number of children in the home, and maternal education. See Table 1 for descriptive statistics and tests of group differences.

Participants were recruited from a five-county region that included both urban and rural areas. Mothers with BPD were recruited from two sources: clinicians in mental health settings and directly from the community. The clinicians included therapists, psychiatrists, nurse practitioners, case managers and other professionals from local hospitals, community outpatient clinics, homeless shelters, professional organizations, and private practice. Clinicians were notified about the study through presentations on BPD, newsletters, and continuing education seminars. They were provided with brochures to hand to their clients. Mothers with BPD were also recruited directly from the community with flyers posted throughout the five-county region. These flyers listed questions about symptoms of BPD and invited those with children age 4–7 to apply. Questions included the following: “Do you fear abandonment in relationships? Do you find it difficult to control your anger? Are you very impulsive? Do your relationships have extreme ups and downs? Have you hurt yourself or threatened to do so?”

Comparison participants were also recruited from two sources: programs for children and directly from the community. Programs included preschools, Head Start and Boys' and Girls' Clubs. Comparison mothers were also recruited directly from the community with flyers. These flyers asked mothers if they had a child aged 4–7 and would like to take part in a study on parent–child interactions.

Procedures and measures

Overall. After a phone screen, two research assistants went to the participant's home or an

Table 1. Child age, child verbal abilities, and demographic differences between the BPD and normative comparison groups

Variable	Whole Sample (<i>N</i> = 60) <i>M</i> (<i>SD</i>)	BPD (<i>n</i> = 30) <i>M</i> (<i>SD</i>)	Comparisons (<i>n</i> = 30) <i>M</i> (<i>SD</i>)	<i>t</i>
Child age (years)	5.33 (0.87)	5.19 (0.84)	5.49 (0.90)	1.35
Verbal ability (PPVT)	102.55 (14.19)	100.67 (16.05)	104.43 (12.04)	1.09
Family yearly income (\$)	32,405 (29,630)	30,902 (20,874)	33,908 (36,684)	0.39
Adults in home	1.82 (0.77)	1.80 (0.71)	1.83 (0.83)	0.17
Children in home	2.47 (1.19)	2.53 (1.31)	2.40 (1.07)	0.43
	%	%	%	χ^2
Child gender (girls)	50	50	50	0.00
Child minority ethnic background	12	10	13	0.16
Mother graduated high school or GED	88	83	93	1.46
Mother has partner	77	77	77	0.00

Note: BPD, borderline personality disorder; PPVT, Peabody Picture Vocabulary Test (Dunn & Dunn, 1997).

alternative meeting place requested by the participant. Informed consent, maternal self-report of BPD symptoms, and demographic information were obtained at this initial visit. If eligible, the participant and her child were then invited to the university for one approximately 3-hr visit. Transportation and babysitting for siblings were provided. At this visit maternal psychopathology was assessed in a clinical interview, and the child was filmed completing stories begun for him or her by an examiner. Individuals who administered the clinical interview were different from those who administered the story-stem completions.

Demographics. Demographic information was collected with a maternal interview (Mt. Hope Family Center, 1995). See Table 1 for details.

Psychiatric diagnoses

BPD. BPD was assessed first on the home visit with a preliminary self-report screen, which was followed up with the Structured Clinical Interview for the *Diagnostic and Statistical Manual of Mental Disorders—Fourth Edition (DSM-IV;* American Psychiatric Association, 1994) Axis II disorders (SCID-II; First, Gibbon, Spitzer, Williams, & Benjamin, 1997).

Current MDD. MDD was also assessed as a control variable with SCID for *DSM-IV* Axis I disorders (SCID-I; First, Gibbon, Spitzer, & Williams, 1996). MDD was only found in the BPD group, where 13% (*n* = 4) mothers currently had MDD.

Borderline features. Borderline features were assessed as part of a self-report measure that assesses psychopathology and personality traits, the Personality Assessment Inventory (PAI; Morey, 1991). The PAI was designed to reflect the multifaceted nature of BPD as revealed in factor analytic studies and has shown high internal consistency in census, college, and clinical samples (Morey, 1991). There are four subscales: *affective instability*, assessing intense and unmodulated emotional experiences especially anger; *identity disturbance* assessing confusion about identity and lack of an integrated sense of self; *negative relationships* assessing acute dependence, fear of abandonment and distrust; and *self-harm*, assessing impulsivity and tendencies to hurt the self when distressed characteristic of individuals with BPD. In the current sample, BPD diagnosis (yes/no) was significantly correlated with *affective instability* ($r = .55, p < .001$), *identity disturbance* ($r = .56, p < .001$), *negative relationships* ($r = .43, p < .01$), and *self-harm* ($r = .59, p < .001$).

Verbal ability. Because narrative creation may depend in part on the child's verbal ability, receptive language was assessed with the Peabody Picture Vocabulary Test, Third Edition (Dunn & Dunn, 1997). Standard scores were calculated for each child.

Narrative story-stem completions. Each child was individually told the beginning of 10 stories, 1 at a time, with the examiner moving figures and props around as though in a play. The figures represented family members and a dog, and props included household items such as a table with dinnerware, a large rock, a bed, and a car. The ethnic background and gender of the child were matched with those of family figures. The examiner presented each story stem about a challenging family situation by moving the figures around and speaking for them with dramatic inflections. The child was then asked by the examiner to complete the stories in the same way: "Show me and tell me what happens now." An initial story about a birthday party, not included in coding or analyses, was used to familiarize the child with the procedure. The narrative stems were always administered in the same order in a session lasting approximately 30 min. All sessions were filmed through a one-way mirror.

The child was asked to complete 10 story stems: 5 story stems designed to activate the attachment system, the attachment story completion task (ASCT; Bretherton, Ridgeway, et al., 1990) and 5 story stems taken from a larger set (which also includes the ASCT) designed in part to assess moral development, the MacArthur Story Stem Battery (Bretherton, Oppenheim, Buchsbaum, Emde, & The MacArthur Narrative Group, 1990).

The five ASCT story stems elicit responses concerning the attachment relationship in increasingly stressful situations: (a) parent as authority figure, in which a child spills his or her juice at dinner and mother disciplines him or her (Spilled Juice); (b) parent as comforter, in which a child falls off a rock and hurts his or her knee (Hurt Knee); (c) parent as protector, in which the child calls for his or her parents at night having seen a monster (Monster in the Bedroom); (d) separation of the child from his or her parents as the parents leave for a

trip, leaving the children with grandma (Departure); and (e) reuniting of child and parents the next day (Reunion).

The five additional MacArthur Story Stem Battery story stems present further challenging situations: (f) parents fighting, in which parents argue over who lost Mom's keys (Lost Keys); (g) an injured child, where the child's hand is burnt when the child takes the pot off the stove to taste the gravy (Hot Gravy); (h) competition between children in which children fight over who gets to ride the toy horse (Horsie); (i) a lost dog, in which the family dog is lost and found (Barney); and (j) a child is left out as the parents send the child to his or her room to play so they can have some time alone, sit on the couch and kiss (Exclusion).

Narrative coding. Narrative emotion coding (Warren, Mantz-Simmons, & Emde, 1993) was utilized to code *role reversal* and *fear of abandonment*. The Child Narrative Coding System (Bickham & Fiese, 1999) was utilized to code *mother-child* and *father-child expectations*. The Narrative Coding Manual, Rochester Version (Robinson, Mantz-Simmons, Macfie, & The MacArthur Narrative Group, 1996) was utilized for the remaining codes. Both child verbalizations and behavioral enactments were included in coding directly from film.

Caregiver-child relationship. Four codes were included: *role reversal* (e.g., child tells fighting parents to: "Stop that! Go to your room!"); *fear of abandonment* (a loss is resolved or attempted to be resolved immediately; e.g., when presented with the Departure story in which mother and father leave the children with grandma for a weekend to go on a trip, the participant insists that the parents change their mind and stay home or take the children with them instead); *mother-child relationship expectations and father-child expectations*. The parent-child expectation codes were coded on 5-point scales. A score of 1 reflects the parent-child relationship is almost always portrayed as dissatisfying, dangerous, and/or unpredictable, with serious and/or willful harm portrayed. A score of 5 reflects the parent-child relationship is consistently portrayed as safe, reliable, rewarding, and fulfilling, and the relationship provides opportunities for success

and satisfaction. Intermediate scores were also given. Three codes were utilized for *self-representations: negative child* (e.g., child hits mom), *incongruent child* (e.g., child cleans up his room then trashes it), and *shameful child* (e.g., child says he is bad).

Emotion regulation. Five codes were included: *reality/fantasy confusion* (e.g., child participant fears that she has killed her parents in reality by killing them in the story), *self/fantasy confusion* (e.g., child participant confuses himself with the fantasy characters in his story and insists that he get a Band-Aid for the child figure, rather than have one of the story characters get it), *fantasy proneness* (e.g., the family floats up into the sky quite unrelated to the story at hand), *intrusion of traumatic material* (e.g., as mom and dad are arguing about who lost mom's car keys, a robber comes into the house and throws everyone into the ocean where they are eaten by alligators), and *narrative coherence*.

Narrative coherence is scored on a 10-point scale based on how well the child confronts the issue raised in the story and how well the child tells the story. A zero denotes failure to respond or "I don't know." A score of 3 denotes that a child exhibits an understanding of the conflict but does not offer any resolution. A score of 5 denotes that a child exhibits an understanding of the conflict but handles it indirectly by offering an easier solution. A score of 7 denotes that child demonstrates understanding of the conflict, offers a coherent resolution without any story embellishment. A score of 10 denotes that a child exhibits a very logical, coherent, sequential series of events that demonstrate an understanding of and resolution to the conflict with a lot of embellishment. Intermediate scores are also given.

Scoring. Presence or absence of the above codes (except narrative coherence and mother-child and father-child relationship expectations) were scored once for each narrative and then summed across the ten narratives, giving a possible minimum of 0 and maximum of 10 for each code. Scores for narrative coherence, which were scored for each narrative, were averaged across the 10 narratives. Scores for mother-child and father-child relationship expectations were each scored once across all 10 narratives.

Composites. In addition to analyzing each code separately, maladaptive composites of the *caregiver-child relationship*, the *self*, and *emotion regulation* were calculated by standardizing each of the individual codes, then summing negatively valenced codes and subtracting positively valenced codes. For example, the maladaptive caregiver-child relationship composite consisted of role reversal plus fear of abandonment minus mother-child relationship expectations minus father-child relationship expectations.

Reliability. Narratives were coded by one of the MacArthur manual's authors. Reliability was established on 25% of the sample with a second coder with an additional 5% of the sample for low base-rate codes. Neither coder was involved in administering the narratives and neither knew the child's group status. Reliability was conducted at the level of each individual narrative rather than summing across narratives. Categorical codes were assessed with kappas to correct for chance agreement (Cohen, 1960): role reversal, $\kappa = .67$ (96% agreement); fear of abandonment, $\kappa = .69$ (96% agreement); negative child, $\kappa = .76$ (94% agreement); incongruent child, $\kappa = .66$ (98% agreement); shameful child, $\kappa = .96$ (99% agreement); reality/fantasy confusion, $\kappa = .81$ (98% agreement); self/fantasy boundary confusion, $\kappa = .76$ (98% agreement); fantasy proneness, $\kappa = .79$ (98% agreement); intrusion of traumatic material, $\kappa = .61$ (94% agreement). Continuous codes were assessed with intraclass correlation coefficients (Winer, Brown, & Michels, 1991): mother-child relationship expectations, $r_i = .74$; father-child relationship expectations, $r_i = .93$; narrative coherence, $r_i = .81$.

Results

Preliminary analyses

There were no significant group differences on child age, child verbal ability, or on demographic variables. See Table 3 for means, standard deviations, and tests of significance. We first conducted two-tailed Pearson bivariate correlations among the narrative variables in the sample as a whole. For *caregiver-child representations*, role reversal, fear of abandonment, mother-child

and father–child relationship expectations were all significantly correlated with each other. For *self-representations*, negative and incongruent child were significantly correlated. For *emotion regulation*, fantasy proneness was significantly correlated both with reality/fantasy confusion and self/fantasy boundary confusion, and narrative coherence was significantly correlated with mother–and father–child relationship expectations and (negatively) with shameful child. See Table 2 for correlation coefficients for narrative variables.

Hypothesis testing

Prior to testing the first three hypotheses, we conducted three overall multivariate analyses of covariance (MANCOVAs) to assess the effect of maternal BPD on caregiver–child relationship representations, self-representations, and emotion regulation, controlling for maternal MDD. To examine the multivariate effects in more detail and to test each hypothesis, we conducted follow-up univariate analyses. In addition to p values, we report observed power and effect size (partial eta squared [η_p^2], the equivalent of ΔR^2 from multiple regression models).

Hypothesis 1. To examine the effect of maternal BPD on caregiver–child relationship representations, we conducted a MANCOVA with maternal BPD status (BPD, no BPD) as the independent variable and MDD as the covariate. Dependent variables included role reversal, fear of abandonment, mother–child relationship expectations, and father–child relationship expectations. There was a significant overall main effect for BPD, Wilks' approximate $F(4, 54) = 3.10$, $\eta_p^2 = .19$, observed power = .78. Univariate F tests for the effect of BPD status revealed significant differences as hypothesized. Children whose mothers had BPD portrayed significantly more role reversal, more fear of abandonment, and more negative mother–child and father–child relationship expectations than did comparisons. See Table 3 for group means, univariate F tests, effect sizes, and observed power.

Hypothesis 2. To examine the effect of maternal BPD on self-representations, we conducted a MANCOVA with maternal BPD status (BPD,

no BPD) as the independent variable and MDD as the covariate. Dependent variables included: negative child, incongruent child, and shameful child. There was a significant overall main effect for BPD, Wilks' approximate $F(3, 55) = 4.65$, $p < .01$, $\eta_p^2 = .20$, observed power = .87. Univariate F tests for the effect of BPD status revealed two out of three significant differences as hypothesized. Children whose mothers had BPD portrayed significantly more incongruent child and shameful child (but not negative child) representations than did comparisons. See Table 3 for group means, univariate F tests, effect sizes, and observed power.

Hypothesis 3. To examine the effect of maternal BPD on emotion regulation, we conducted a MANCOVA with maternal BPD status (BPD, no BPD) as the independent variable and MDD as the covariate. Dependent variables included reality/fantasy confusion, self/fantasy boundary confusion, fantasy proneness, intrusion of traumatic material, and narrative coherence. There was a significant overall effect for BPD, Wilks' approximate $F(5, 53) = 3.76$, $p < .01$, $\eta_p^2 = .26$, observed power = .91. Univariate F tests for BPD status revealed significant differences as hypothesized. Children whose mothers had BPD demonstrated more reality/fantasy confusion, more self/fantasy boundary confusion, more fantasy proneness, lower narrative coherence, and marginally more intrusion of traumatic material than did comparisons. See Table 3 for group means, univariate F tests, effect sizes, and observed power.

To test the remaining hypotheses, we conducted two-tailed Pearson bivariate correlations.

Hypothesis 4. As hypothesized, the child's maladaptive child–caregiver relationship composite was significantly correlated with maternal identity disturbance and self-harm, but, contrary to hypothesis, not with maternal negative relationships.

Hypothesis 5. As hypothesized, the child's maladaptive self-composite was significantly correlated with maternal self-harm.

Hypothesis 6. As hypothesized, the child's emotion regulation composite was significantly correlated with maternal identity disturbance,

Table 2. Intercorrelations among narrative variables in the whole sample ($N = 60$)

	1	2	3	4	5	6	7	8	9	10	11
1. Role reversal	—										
2. Fear of abandonment	.36**	—									
3. Mother-child relationship expectations	-.27*	-.40**	—								
4. Father-child relationship expectations	-.32*	-.24	.83***	—							
5. Negative child	.22	.34**	-.47***	-.45***	—						
6. Incongruent child	.16	.42**	-.21	-.22	.47***	—					
7. Shameful child	.29*	.18	-.11	-.10	-.15	-.10	—				
8. Reality/fantasy confusion	.13	.20	-.19	-.26*	.29*	.20	-.03	—			
9. Self/fantasy confusion	.44***	.34**	-.19	-.22	.16	.29*	.11	.23	—		
10. Fantasy proneness	.04	.11	-.08	-.19	.19	.26*	-.09	.57***	.35**	—	
11. Intrusion of traumatic material	.24	.08	-.27*	-.35**	.36**	.32*	-.11	.45***	.19	.52***	—
12. Narrative coherence	-.05	-.08	.53***	.47**	-.17	-.17	-.35**	.06	-.01	.08	-.06

* $p < .05$. ** $p < .01$. *** $p < .001$.

Table 3. Group differences in narrative variables: Univariate *F* tests, effect size, and observed power

Narrative Variable	Whole Sample (<i>N</i> = 60) <i>M</i> (<i>SD</i>)	BPD (<i>n</i> = 30) <i>M</i> (<i>SD</i>)	Comparisons (<i>n</i> = 30) <i>M</i> (<i>SD</i>)	<i>F</i> (<i>df</i>)	η^2	Observed Power
Caregiver-child relation. represent.				<i>F</i> (1, 57)		
Role reversal	0.62 (0.83)	0.87 (1.01)	0.37 (0.49)	4.32*	.07	.53
Fear of abandonment	0.57 (0.74)	0.87 (0.77)	0.27 (0.58)	8.72**	.13	.83
Mother-child relation. expect.	2.93 (1.48)	2.43 (1.33)	3.43 (1.48)	5.65*	.09	.65
Father-child relation. expect.	3.27 (1.42)	2.83 (1.42)	3.70 (1.32)	4.48*	.07	.55
Self-representations				<i>F</i> (1, 57)		
Negative child	1.37 (1.52)	1.70 (1.80)	1.03 (1.10)	0.89	.02	.15
Incongruent child	0.10 (0.30)	0.20 (0.41)	0.00 (0.00)	4.29*	.07	.53
Shameful child	0.08 (0.28)	0.17 (0.38)	0.00 (0.00)	7.27**	.11	.76
Emotion regulation				<i>F</i> (1, 57)		
Reality/fantasy confusion	0.27 (0.66)	0.43 (0.82)	0.10 (0.40)	4.34*	.07	.54
Self/fantasy confusion	0.20 (0.55)	0.40 (0.72)	0.00 (0.00)	7.75**	.12	.78
Fantasy proneness	0.60 (1.40)	1.03 (1.87)	0.17 (0.38)	5.74*	.09	.65
Intrusion of traumatic material	1.20 (1.71)	1.57 (1.98)	0.83 (1.32)	2.98†	.05	.40
Narrative coherence	6.17 (1.91)	5.58 (2.06)	6.75 (1.57)	4.83*	.08	.58

†*p* < .10. **p* < .05. ***p* < .01.

negative relationships, and self-harm, and marginally significantly correlated with maternal affective instability. See Table 4 for correlation coefficients.

Discussion

We assessed representations of the caregiver-child relationship and of the self, and emotion regulation in children age 4-7 whose mothers have BPD versus normative comparisons using a narrative story-stem completion measure. The study extended developmental research on offspring of mothers with BPD from infancy to early childhood, from the level of sensorimotor functioning to the level of representation. Maladaptive representations and poor emotion regulation may increase a child's own likelihood of developing BPD in early adulthood.

Findings suggest problems for the children whose mothers have BPD, controlling for maternal MDD. First, children whose mothers have BPD told stories in which the caregiver-child relationship was characterized by more role rever-

sal, more fear of abandonment, and more negative mother-child and father-child relationship expectations. Second, children whose mothers have BPD told stories in which the child's self was represented as more incongruent and shameful. Contrary to hypothesis, representations of the self were not more negative, however. Negative representations have been associated with the experience of maltreatment (Toth, Cicchetti, Macfie, & Emde, 1997; Toth, Cicchetti, Macfie, Maughan, et al., 2000). Although many adults with BPD report having been maltreated themselves (Zanarini, 2000) not all will maltreat their own children (Egeland, Jacobvitz, & Sroufe, 1988). Third, children whose mothers have BPD demonstrated impaired emotion regulation. They were less able to maintain a distinction between fantasy and reality and between self and fantasy, more likely to veer away from the issue presented into fantasy proneness, were less coherent, and were marginally more likely to bring intrusive traumatic themes into their stories.

Fourth, in the sample as a whole, maladaptive composites of the child's caregiver-child

Table 4. Correlations between child maladaptive narrative composites and maternal borderline features (PAI) in the sample as a whole ($N = 60$)

Maternal Borderline Features (PAI)	Child Maladaptive Narrative Composites		
	Child–Caregiver Relation. Represent.	Self-Represent.	Emotion Regul.
Affective instability	.19	.10	.24†
Identity disturbance	.30*	.20	.42**
Negative relationships	.18	.21†	.35**
Self-harm	.26*	.26*	.25*

Note: PAI, Personality Assessment Inventory (Morey, 1991).

† $p \leq .10$. * $p < .05$. ** $p < .01$.

relationship representations, the child's self-representations and the child's emotion regulation were each significantly correlated with maternal borderline features. Specifically, the child's maladaptive caregiver–child relationship composite was associated with his or her mother's confusion about her identity and her self-harming behaviors. Moreover, a child's maladaptive self-composite was also associated with his or her mother's self-harming behaviors. Furthermore, a child's maladaptive emotion regulation composite was associated with his or her mother's confusion about her identity, negative relationships, self-harming behavior, and was marginally significantly associated with affective instability.

Putative precursors to BPD

Do results suggest that young children whose mothers have BPD are at risk of developing BPD themselves in early adulthood? Not all or even most children of mothers with BPD will develop BPD. Indeed, only 11.5% of first-degree relatives of individuals with BPD also have BPD (Nigg & Goldsmith, 1994). However, exposure to risk factors associated with BPD: a vulnerable temperament, and failure at tasks of attachment, self-development, and emotion regulation may increase the likelihood of developing BPD.

Adults with BPD do not report histories of being in treatment as children (Zanarini, Frankenburg, Khera, & Bleichmar, 2001), so it is important to examine possible precursors to BPD that do not necessarily imply psychopathology in childhood but may develop into psychopathology

in early adulthood. Heterotypic continuity may underlie development such that the same construct may look dissimilar in different developmental periods but carry the same meaning from the past into the present via internal working models (Carlson et al., 2004).

Thus, a young child who demonstrates fear of abandonment in his or her narratives may grow up to demonstrate the same fear of abandonment as a young adult in relationships. This same young adult may still cling to transitional objects such as teddy bears on arrival at the hospital diagnosed with BPD (Gunderson, 2001). In addition, a young child who demonstrates negative expectations of both mother–child and father–child relationships in his or her narratives may carry both anger and mistrust about others into new relationships and later develop intense, unstable relationships marked by inappropriate angry outbursts as an adult. Furthermore, a young child who demonstrates a shameful and incongruent self in his or her narratives may grow up to use self-injurious behavior as a way to temporarily relieve feelings of shame and create a unified sense of self (Yates, 2004). Moreover, a young child who demonstrates poor emotion regulation in his or her narratives may grow up to experience the same inability to draw upon an internalized self-soothing process as a young adult in times of emotional distress (Adler & Buie, 1979) and turn instead to drugs, alcohol, or food to soothe the self (Bradley & Westen, 2005).

Findings also include associations between maternal borderline features and maladaptive narrative composites. Importantly, a mother's

self-harming behavior was associated with her child's maladaptive relationship, self, and emotion regulation composites. This is particularly significant because pathways to self-harming behavior are thought to include failure to experience relationships as reliable or rewarding, the self to be worthy of care, and failure to modulate emotion and arousal. Failure with stage-salient issues of attachment, self, and self-regulation may then lead to the development of compensatory maladaptive strategies including self-harm to find a way to continue to address future stage-salient issues in adolescence (Yates, 2004). Children whose mothers have BPD may therefore be at high risk of developing self-harming behaviors in early adulthood.

Emotion regulation, dissociation, and disorganized attachment

Problems with emotion regulation found in the current study overlap with those reflecting dissociation in maltreated children (Macfie et al., 2001). Dissociation, which refers to disruptions in the normal integration of memories, perception, and identity, is a symptom of BPD (American Psychiatric Association, 1994). Dissociation has been conceptualized as a failure of information processing in the face of trauma, a breakdown in emotion regulation because of flooding of the sympathetic nervous system with stress hormones and neurotransmitters (Bower & Sivers, 1998). One pathway to dissociation is from disorganized attachment (Carlson, 1998; Ogawa, Sroufe, Weinfield, Carlson, & Egeland, 1997), and most infants whose mothers have BPD are classified as disorganized (Hobson et al., 2005). Moreover, children disorganized as infants become more vulnerable to posttraumatic stress disorder in childhood (MacDonald et al., 2008). This difficulty with emotion regulation in the face of stress may also have implications for the development of BPD. Indeed, dissociation mediates the relationship between the experience of childhood sexual abuse and self-harming behavior in adults (Yates, Carlson, & Egeland, 2008). Furthermore, children whose mothers have BPD may be at risk of sexual abuse because many women with BPD report sexual abuse as children (Zanarini et al., 1997).

There are three known pathways to disorganized attachment: the experience of maltreatment

(Carlson, Cicchetti, Barnett, & Braunwald, 1989), mothers' unresolved experience of loss or trauma (Main & Hesse, 1990), and mothers' disrupted affective communication that stems from hostile/helpless states of mind (Lyons-Ruth, Bronfman, & Atwood, 1999; Lyons-Ruth, Melnick, Patrick, & Hobson, 2007; Lyons-Ruth, Yellin, Melnick, & Atwood, 2005). Each pathway may be relevant for infants whose mothers have BPD. Disorganized attachment is associated with negative expectations of others, a negative sense of self, and disrupted emotion regulation. Thus, disorganized attachment may be a necessary but not sufficient precursor for the development of BPD, an inability to handle stress underlying both, leading to faulty emotion regulation including self-harming behavior.

Preventive interventions

Given the size of the public health problem presented by BPD, research to inform prevention is warranted. In addition to being responsible for chronically high levels of individual distress, individuals with BPD impose a severe burden on the health care system. Individuals with BPD utilize mental health services at higher rates than does any diagnostic group other than schizophrenia (Swartz et al., 1990). Seventy to 90% of individuals with BPD repeatedly attempt suicide or make suicidal gestures (Gunderson & Ridolfi, 2001; Linehan & Heard, 1999), which involve intensive utilization of mental health services (Roy, 2001). Although great strides have been made in treatment (Bateman & Fonagy, 1999, 2001, 2008; Gunderson, 2001; Levy et al., 2006; Linehan, 1993), BPD remains a disorder from which many mental health practitioners recoil. More needs to be known about the development of BPD to inform preventive interventions. Results of the current study suggest the importance of interventions with children whose mothers have BPD: interventions that are designed to alter representational models of the caregiver-child relationship and of the self, and improve emotion regulation.

Dyadic child-parent psychotherapy (Fraiberg, Adelson, & Shapiro, 1975; Lieberman, 1992) is an attachment-based therapy that has been successful in altering representational models. A mother and her young child (infant, toddler, or

preschooler) meet together with the therapist. The mother feels understood by the therapist and learns more about her own and her child's feelings, beliefs, and needs, so that the mother-child relationship becomes a greater source of security to the child. Indeed, child-parent psychotherapy leads to an increase in attachment security in depressed mother-toddler pairs (Cicchetti et al., 1999; Toth et al., 2006) and an increase in positive, and a decrease in negative, representations in maltreated children's stories (Toth et al., 2002). Both a mother with BPD and her child may benefit, and development for both may return to a more adaptive pathway. Dyadic therapy may also help mother and child construct a narrative that is coherent about traumatic experiences, the coherence both reflecting and contributing to emotion regulation skills (Oppenheim, Nir, et al., 1997). Interestingly, an attachment-based intervention that focuses on the caregiver of children in foster care has been shown to improve the child's ability to cope with stress at the level of cortisol in the hypothalamus-pituitary-adrenal axis (Dozier, Peloso, Lewis, Laurenceau, & Levine, 2008). Moreover, attachment mediated the relationship between maternal symptoms of depression and narrative representations (Toth et al., 2009). Interventions designed to improve attachment in the caregiver-child relationship may be warranted for young children whose mothers have BPD.

Strengths and Limitations

The study's strengths include filmed and reliably coded observational data of children rather than relying on adult observer report measures. Moreover, larger numbers of children were sampled in the same developmental period compared with previous studies of children whose mothers have BPD. Furthermore, in addition to categorical diagnoses of BPD assessed in a clinical interview,

continua of borderline features were assessed in a self-report measure. Categorical diagnoses are meaningful to clinicians, whereas symptoms on a continuum have more statistical power and they can be assessed for all participants. In addition, because mothers with BPD were recruited both from clinician referrals and directly from the community, the sample has greater generalizability to the population as a whole than it would if referred solely from women in treatment. Furthermore, mothers were recruited from low SES backgrounds. Because BPD is more often found in individuals from low SES backgrounds, this also increases generalizability to the population as a whole (Cohen et al., 2008; Grant et al., 2008).

Limitations of the current study include the fact that risk factors for the development of BPD such as the experience of maltreatment, loss of caregiver, and temperamental factors were not assessed. Also, MDD was controlled for statistically. It would be better to have a second comparison group of depressed mothers and their children in addition to a normative comparison group.

Conclusion

A developmental psychopathology perspective makes it possible to study development in an at-risk sample and learn more about not only psychopathology but also about normal development. Children whose mothers have BPD shed light on representational development and the development of emotion regulation in all children. Moreover, the usefulness of the narrative story-stem completion measure is extended to offspring of women diagnosed with BPD. Finally, following these children over time will help us to better understand problems with early stage-salient issues of attachment, self-development and emotion regulation, and under what circumstances such problems lead to BPD.

References

- Adler, G., & Buie, D. (1979). Aloneness and borderline psychopathology: The possible relevance of child development issues. *International Journal of Psychoanalysis*, 60, 83-96.
- Ainsworth, M. D. S., Blehar, M. C., Waters, E., & Wall, S. (1978). *Patterns of attachment: A psychological study of the strange situation*. Hillsdale, NJ: Erlbaum.
- American Psychiatric Association. (1994). *Diagnostic and statistical manual of mental disorders* (4th ed.). Washington, DC: Author.
- Bar-Haim, Y., Fox, N. A., VanMeenen, K. M., & Marshall, P. J. (2004). Children's narratives and patterns of cardiac reactivity. *Developmental Psychobiology*, 44, 238-249.

- Barnow, S., Spitzer, C., Grabe, H. J., Kessler, C., & Freyberger, H. J. (2006). Individual characteristics, familial experience, and psychopathology in children of mothers with borderline personality disorder. *Journal of the American Academy of Child & Adolescent Psychiatry*, *65*, 965–972.
- Bateman, A., & Fonagy, P. (1999). Effectiveness of partial hospitalization in the treatment of borderline personality disorder: A randomized controlled trial. *American Journal of Psychiatry*, *156*, 1563–1569.
- Bateman, A., & Fonagy, P. (2001). Treatment of borderline personality disorder with psychoanalytically oriented partial hospitalization: An 18-month follow-up. *American Journal of Psychiatry*, *158*, 36–42.
- Bateman, A., & Fonagy, P. (2008). 8-Year follow-up of patients treated for borderline personality disorder: Mentalization-based treatment versus treatment as usual. *American Journal of Psychiatry*, *165*, 631–638.
- Belden, A., Sullivan, J. P., & Luby, J. L. (2007). Depressed and healthy preschoolers' internal representations of their mothers' caregiving: Associations with observed caregiving behaviors one year later. *Attachment and Human Development*, *9*, 239–254.
- Bezirgianian, S., Cohen, P., & Brook, J. S. (1993). The impact of mother-child interaction on the development of borderline personality disorder. *American Journal of Psychiatry*, *150*, 1836–1842.
- Bickham, N., & Fiese, B. (1999). *Child Narrative Coding System*. Unpublished manuscript, Syracuse University.
- Bower, G. H., & Sivers, H. (1998). Cognitive impact of traumatic events. *Development and Psychopathology*, *10*, 625–654.
- Bowlby, J. (1982). *Attachment and loss. Vol. 1. Attachment*. New York: Basic Books. (Original work published 1969)
- Bowlby, J. (1988). *A secure base*. New York: Basic Books.
- Bradley, R., & Westen, D. (2005). The psychodynamics of borderline personality disorder: A view from developmental psychopathology. *Development and Psychopathology*, *17*, 927–957.
- Bretherton, I., & Munholland, K. A. (2008). Internal working models in attachment relationships: Elaborating a central construct in attachment theory. In J. Cassidy & P. R. Shaver (Eds.), *Handbook of attachment: Theory, research and clinical applications* (2nd ed.). New York: Guilford Press.
- Bretherton, I., Oppenheim, D., Buchsbaum, H., Emde, R. N., & The MacArthur Narrative Group. (1990). *MacArthur Story Stem Battery (MSSB)*. Unpublished manuscript.
- Bretherton, I., Ridgeway, D., & Cassidy, J. (1990). Assessing internal working models of the attachment relationship. In M. T. Greenberg, D. Cicchetti, & M. Cummings (Eds.), *Attachment in the preschool years* (pp. 273–308). Chicago: University of Chicago Press.
- Carlson, E. A. (1998). A prospective longitudinal study of attachment disorganization/disorientation. *Child Development*, *69*, 1107–1128.
- Carlson, E. A., Sroufe, L. A., & Egeland, B. (2004). The construction of experience: A longitudinal study of representations and behavior. *Child Development*, *75*, 66–83.
- Carlson, V., Cicchetti, D., Barnett, D., & Braunwald, K. (1989). Finding order in disorganization: Lessons from research on maltreated infants' attachments to their caregivers. In D. Cicchetti & V. Carlson (Eds.), *Child maltreatment: Theory and research on the causes and consequences of child abuse and neglect* (pp. 494–528). New York: Cambridge University Press.
- Cassidy, J. (1988). Child-mother attachment and the self in six-year-olds. *Child Development*, *59*, 121–134.
- Cicchetti, D. (1984). The emergence of developmental psychopathology. *Child Development*, *55*, 1–7.
- Cicchetti, D., & Hinshaw, S. P. (2002). Editorial. Prevention and intervention science: Contributions to developmental theory. *Development and Psychopathology*, *14*, 667–672.
- Cicchetti, D., Rogosch, R., & Toth, S. L. (2000). The efficacy of toddler-parent psychotherapy for fostering cognitive development in offspring of depressed mothers. *Journal of Abnormal Child Psychology*, *28*, 135–148.
- Cicchetti, D., & Toth, S. L. (1992). The role of developmental theory in prevention and intervention. *Development and Psychopathology*, *4*, 489–493.
- Cicchetti, D., & Toth, S. L. (1998). The development of depression in children and adolescents. *American Psychologist*, *53*, 221–241.
- Cicchetti, D., & Toth, S. L. (2006). Developmental psychopathology and preventive intervention. In K. A. Renninger & I. E. Sigel (Eds.), *Handbook of child psychology*. Hoboken, NJ: Wiley.
- Cicchetti, D., Toth, S. L., & Rogosch, F. (1999). The efficacy of toddler-parent psychotherapy to increase attachment security in offspring of depressed mothers. *Attachment and Human Development*, *1*, 34–66.
- Cohen, J. (1960). A coefficient of agreement for nominal scales. *Education and Psychological Measurement*, *20*, 37–46.
- Cohen, P., Chen, H., Gordon, K., Johnson, J., Brook, J., & Kasen, S. (2008). Socioeconomic background and the developmental course of schizotypal and borderline personality disorder symptoms. *Development and Psychopathology*, *20*, 633–650.
- Crandell, L. E., Patrick, M., & Hobson, R. P. (2003). "Still-face" interactions between mothers with borderline personality disorder and their 2-month-old infants. *British Journal of Psychiatry*, *183*, 239–247.
- Crick, N. R., Murray-Close, D., & Woods, K. (2005). Borderline personality features in childhood: A short-term longitudinal study. *Development and Psychopathology*, *17*, 1051–1070.
- Curtis, W. J., & Cicchetti, D. (2007). Emotion and resilience: A multi-level investigation of hemispheric electroencephalogram asymmetry and emotion regulation in maltreated and nonmaltreated children. *Development and Psychopathology*, *19*, 811–840.
- Downey, G., & Coyne, J. C. (1990). Children of depressed parents: An integrative review. *Psychological Bulletin*, *108*, 50–76.
- Dozier, M., Peloso, E., Lewis, E., Laurenceau, J., & Levine, S. (2008). Effects of an attachment-based intervention on the cortisol production of infants and toddlers in foster care. *Development and Psychopathology*, *20*, 845–859.
- Dunn, L. M., & Dunn, L. (1997). *Peabody Picture Vocabulary Test* (3rd ed.). Circle Pines, MN: American Guidance Service.
- Egeland, B. R., Jacobvitz, D., & Sroufe, L. A. (1988). Breaking the cycle of abuse. *Child Development*, *58*, 1080–1088.
- Feldman, R. B., Zerkowicz, P., Weiss, M., Vogel, J., Heyman, M., & Paris, J. (1995). A comparison of the families of mothers with borderline and nonborderline personality disorders. *Comprehensive Psychiatry*, *36*, 157–163.
- First, M. B., Gibbon, M., Spitzer, R. L., & Williams, J. B. W. (1996). *User's guide for the Structured Clinical Interview for DSM-IV Axis I Disorders—Research version*. Washington, DC: American Psychiatric Press.

- First, M. B., Gibbon, M., Spitzer, R. L., Williams, J. B. W., & Benjamin, L. S. (1997). *Structured Clinical Interview for DSM-IV Axis II personality disorders: SCID-II*. Washington, DC: American Psychiatric Press.
- Fonagy, P., Target, M., & Gergely, G. (2000). Attachment and borderline personality disorder. *Psychiatric Clinics of North America*, 23, 103–122.
- Fraiberg, S., Adelson, E., & Shapiro, V. (1975). Ghosts in the nursery. *Journal of the American Academy of Child & Adolescent Psychiatry*, 14, 387–421.
- Grant, B. F., Chou, S. P., Goldstein, R. B., Huang, B., Stinson, F. S., Saha, T. D., et al. (2008). Prevalence, correlates, disability, and comorbidity of DSM-IV borderline personality disorder: Results from the Wave 2 National Epidemiologic Survey on Alcohol and Related Conditions. *Journal of Clinical Psychiatry*, 69, 533–545.
- Green, J., Stanley, C., & Peters, S. (2007). Disorganized attachment representation and atypical parenting in young school age children with externalizing disorder. *Attachment and Human Development*, 9, 207–222.
- Grych, J. H., Wachsmuth-Schlaefler, T., & Klockow, L. L. (2002). Interparental aggression and young children's representations of family relationships. *Journal of Family Psychology*, 16, 259–272.
- Gunderson, J. G. (1996). The borderline patient's intolerance of aloneness: Insecure attachments and therapist availability. *American Journal of Psychiatry*, 153, 752–758.
- Gunderson, J. G. (2001). *Borderline personality disorder: A clinical guide*. Washington, DC: American Psychiatric Publishing.
- Gunderson, J. G., Carpenter, W., & Strauss, J. (1975). Borderline and schizophrenic patients: A comparative study. *American Journal of Psychiatry*, 132, 1257–1264.
- Gunderson, J. G., & Ridolfi, M. A. (2001). Borderline personality disorder: Suicidality and self-mutilation. *Annals of the New York Academy of Sciences*, 932, 60–83.
- Heard, H. L., & Linehan, M. M. (1993). Problems of self and borderline personality disorder: A dialectical behavioral analysis. In Z. V. Segal & S. J. Blatt (Eds.), *The self in emotional distress: Cognitive and psychodynamic perspectives* (pp. 301–333). New York: Guilford Press.
- Herman, J. L., Perry, J. C., & Kolk, B. A. v. d. (1989). Childhood trauma in borderline personality disorder. *American Journal of Psychiatry*, 146, 490–495.
- Hesse, E., & Main, M. (2006). Frightened, threatening, and dissociative parental behavior in low-risk samples: Description, discussion, and interpretations. *Development and Psychopathology*, 18, 309–343.
- Hill, J., Fonagy, P., Lancaster, G., & Broyden, N. (2007). Aggression and intentionality in narrative responses to conflict and distress story stems: An investigation of boys with disruptive behavior problems. *Attachment and Human Development*, 9, 223–237.
- Hobson, R. P., Patrick, M., Crandell, L., Garcia-Perez, R., & Lee, A. (2005). Personal relatedness and attachment in infants of mothers with borderline personality disorder. *Development and Psychopathology*, 17, 329–347.
- Laporte, L., & Guttman, H. (1996). Traumatic childhood experiences as risk factors for borderline and other personality disorders. *Journal of Personality Disorders*, 10, 247–259.
- Lenzenweger, M. F., & Cicchetti, D. (2005). Toward a developmental psychopathology approach to borderline personality disorder. *Development and Psychopathology*, 17, 893–898.
- Levy, K. N., Meehan, K. B., Kelly, K. M., Reynoso, J. S., Weber, M., Clarkin, J. F., et al. (2006). Change in attachment patterns and reflective functioning in a randomized control trial of transference-focused psychotherapy for borderline personality disorder. *Journal of Consulting and Clinical Psychology*, 74, 1027–1040.
- Lieberman, A. F. (1992). Infant–parent psychotherapy with toddlers. *Development and Psychopathology*, 4, 559–574.
- Linehan, M. M. (1993). *Cognitive-behavioral treatment of borderline personality disorder*. New York: Guilford Press.
- Linehan, M. M., & Heard, H. L. (1999). Borderline personality disorder: Costs, course, and treatment outcomes. In N. E. Miller & K. M. Magruder (Eds.), *Cost-effectiveness of psychotherapy: A guide for practitioners, researchers, and policymakers* (pp. 291–305). New York: Oxford University Press.
- Liotti, G., & Pasquini, P. (2000). Predictive factors for borderline personality disorder: Patients' early traumatic experiences and losses suffered by the attachment figure. *Acta Psychiatrica Scandinavica*, 102, 282–289.
- Ludolph, P. S., Westen, D., Mislis, B., Jackson, A., Wixom, J., & Wiss, F. C. (1990). The borderline diagnosis in adolescents: Symptoms and developmental history. *American Journal of Psychiatry*, 147, 470–476.
- Lyons-Ruth, K., Bronfman, E., & Atwood, G. (1999). A relational diathesis model of hostile–helpless states of mind. In J. Solomon & C. George (Eds.), *Attachment disorganization* (pp. 33–69). New York: Guilford Press.
- Lyons-Ruth, K., Melnick, S., Patrick, M., & Hobson, R. P. (2007). A controlled study of hostile–helpless states of mind among borderline and dysthymic women. *Attachment and Human Development*, 9, 1–16.
- Lyons-Ruth, K., Yellin, C., Melnick, S., & Atwood, G. (2005). Expanding the concept of unresolved mental states: Hostile/helpless states of mind on the Adult Attachment Interview are associated with disrupted mother–infant communication and infant disorganization. *Development and Psychopathology*, 17, 1–23.
- MacDonald, H. Z., Beeghly, M., Grant-Knight, W., Augustyn, M., Woods, R. W., Cabral, H., et al. (2008). Longitudinal association between infant disorganized attachment and childhood posttraumatic stress symptoms. *Development and Psychopathology*, 20, 493–508.
- Macfie, J. (2009). Development in children and adolescents whose mothers have borderline personality disorder. *Child Development Perspectives*, 3, 66–71.
- Macfie, J., Cicchetti, D., & Toth, S. L. (2001). The development of dissociation in maltreated preschool-aged children. *Development and Psychopathology*, 13, 233–253.
- Macfie, J., Fitzpatrick, K. L., Rivas, E. M., & Cox, M. J. (2008). Independent influences on mother–toddler role reversal: Infant–mother attachment disorganization and role reversal in mother's childhood. *Attachment and Human Development*, 10, 29–39.
- Macfie, J., Houts, R. M., McElwain, N. L., & Cox, M. J. (2005). The effect of father–toddler and mother–toddler role reversal on the development of behavior problems in kindergarten. *Social Development*, 14, 514–531.
- Macfie, J., McElwain, N. L., Houts, R. M., & Cox, M. J. (2005). Intergenerational transmission of role reversal between parent and child: Dyadic and family systems internal working models. *Attachment and Human Development*, 7, 51–65.
- Macfie, J., Toth, S. L., Rogosch, F. A., Robinson, J. L., Emde, R. N., & Cicchetti, D. (1999). Effect of maltreatment on preschoolers' narrative representations of responses to relieve distress and of role reversal. *Developmental Psychology*, 35, 460–465.

- Main, M., & Hesse, E. (1990). Parents' unresolved traumatic experiences are related to infant disorganized attachment status: Is frightened and/or frightening parental behavior the linking mechanism? In M. T. Greenberg, D. Cicchetti, & E. M. Cummings (Eds.), *Attachment in the preschool years: Theory, research, and intervention* (pp. 161–182). Chicago: University of Chicago Press.
- Main, M., Kaplan, N., & Cassidy, J. C. (1985). Security in infancy, childhood, and adulthood: A move to the level of representation. *Monographs of the Society for Research in Child Development*, 50(1–2, Serial No. 209), 66–104.
- Main, M., & Solomon, J. (Eds.). (1990). *Procedures for identifying infants as disorganized/disoriented during the Ainsworth strange situation*. Chicago: University of Chicago Press.
- Mednick, S. A., & McNeil, T. F. (1968). Current methodology in research on the etiology of schizophrenia: Serious difficulties which suggest the use of the high-risk-group method. *Psychological Bulletin*, 70, 681–693.
- Mt. Hope Family Center. (1995). *Mt. Hope Demographic Interview*. Unpublished manuscript, University of Rochester.
- Morey, L. C. (1991). *Personality Assessment Inventory*. Lutz, FL: Psychological Assessment Resources.
- Newman, L. K., Stevenson, C. S., Bergman, L. R., & Boyce, P. (2007). Borderline personality disorder, mother–infant interaction and parenting perceptions: Preliminary findings. *Australian and New Zealand Journal of Psychiatry*, 41, 598–605.
- Nigg, J. T., & Goldsmith, H. H. (1994). Genetics of personality disorders: Perspectives from personality and psychopathology research. *Psychological Bulletin*, 115, 346–380.
- Nigg, J. T., Lohr, N. E., Westen, D., Gold, L. J., & Silk, K. R. (1992). Malevolent object representations in borderline personality disorder and major depression. *Journal of Abnormal Psychology*, 101, 61–67.
- Ogawa, J. R., Sroufe, L. A., Weinfield, N. S., Carlson, E. A., & Egeland, B. (1997). Development and the fragmented self: A longitudinal study of dissociative symptomatology in a normative sample. *Development and Psychopathology*, 9, 855–879.
- Oppenheim, D. (2006). Child, parent, and parent–child emotion narratives: Implications for developmental psychopathology. *Development and Psychopathology*, 18, 771–790.
- Oppenheim, D., Emde, R. N., & Warren, S. L. (1997). Children's narrative representations of mothers: Their development and associations with child and mother adaptation. *Child Development*, 68, 127–138.
- Oppenheim, D., Nir, A., Warren, S. L., & Emde, R. N. (1997). Emotion regulation in mother–child narrative co-construction: Associations with children's narratives and adaptation. *Developmental Psychology*, 33, 284–294.
- Oppenheim, D., & Waters, H. (1995). Narrative processes and attachment representations: Issues of development and assessment. *Monographs of the Society for Research in Child Development*, 60, 197–215.
- Paris, J. (1993). Long-term outcome of borderline personality disorder: Implications for treatment. *Canadian Journal of Psychiatry*, 37, 28–34.
- Posner, M. I., Rothbart, M. K., Vizueta, N., Thomas, K. M., Levy, K. N., Fossella, J., et al. (2003). An approach to the psychobiology of personality disorders. *Development and Psychopathology*, 15, 1093–1106.
- Putnam, K. M., & Silk, K. R. (2005). Emotion dysregulation and the development of borderline personality disorder. *Development and Psychopathology*, 17, 899–925.
- Robinson, J. L., Mantz-Simmons, L., Macfie, J., & the MacArthur Narrative Group. (1996). *The narrative coding manual, Rochester version*. Unpublished manuscript.
- Rogosch, F. A., & Cicchetti, D. (2005). Child maltreatment, attention networks, and potential precursors to borderline personality disorder. *Development and Psychopathology*, 17, 1071–1089.
- Roy, A. (2001). Consumers of mental health services. *Suicide and Life Threatening Behavior*, 31, 60–83.
- Sameroff, A. J., & Chandler, M. J. (1975). Reproductive risk and the continuum of caretaking causality. In F. D. Horowitz (Ed.), *Review of child development research* (Vol. 4, pp. 187–244). Chicago: University of Chicago Press.
- Scarlett, W. G., & Wolf, D. (1979). When it's only make-believe: The construction of a boundary between fantasy and reality in storytelling. *New Directions for Child Development*, 6, 29–40.
- Schechter, D. S., Zygmont, A., Coates, S. W., Davies, M., Trabka, K. A., McCaw, J., et al. (2007). Caregiver traumatization adversely impacts young children's mental representations on the MacArthur Story Stem Battery. *Attachment and Human Development*, 9, 187–205.
- Shields, A., Ryan, R. M., & Cicchetti, D. (2001). Narrative representations of caregivers and emotion dysregulation as predictors of maltreated children's rejection by peers. *Developmental Psychology*, 37, 321–337.
- Siever, L. J., & Davis, K. L. (1991). A psychobiological perspective on the personality disorders. *American Journal of Psychiatry*, 148, 1647–1658.
- Solomon, J., George, C., & DeJong, A. (1995). Children classified as controlling at age six: Evidence of disorganized strategies and aggression at home and at school. *Development and Psychopathology*, 7, 447–463.
- Sroufe, L. A., Egeland, B., Carlson, E. A., & Collins, W. A. (2005). *The development of the person: The Minnesota Study of Risk and Adaptation From Birth to Adulthood*. New York: Guilford Press.
- Sroufe, L. A., & Rutter, M. (1984). The domain of developmental psychopathology. *Child Development*, 55, 17–29.
- Stadelmann, S., Perren, S., Wyl, A. v., & Klitzing, K. v. (2007). Associations between family relationships and symptoms/strengths at kindergarten age: What is the role of children's parental representations? *Journal of Child Psychology and Psychiatry*, 48, 996–1004.
- Stone, M. H. (1990). *The fate of borderline patients*. New York: Guilford Press.
- Swartz, M., Blazer, D., George, L., & Winfield, I. (1990). Estimating the prevalence of borderline personality disorder in the community. *Journal of Personality Disorders*, 4, 257–272.
- Torgersen, S., Lygren, S., Oien, P. A., Skre, I., Onstad, S., Edvardsen, J., et al. (2000). A twin study of personality disorders. *Comprehensive Psychiatry*, 41, 416–425.
- Toth, S. L., Cicchetti, D., Macfie, J., & Emde, R. N. (1997). Representations of self and other in the narratives of neglected, physically abused, and sexually abused preschoolers. *Development and Psychopathology*, 9, 781–796.
- Toth, S. L., Cicchetti, D., Macfie, J., Maughan, A., & VanMeenen, K. (2000). Narrative representations of caregivers and self in maltreated preschoolers. *Attachment and Human Development*, 2, 271–305.
- Toth, S. L., Cicchetti, D., Macfie, J., Rogosch, F. A., & Maughan, A. (2000). Narrative representations of moral affiliative and conflictual themes and behavioral problems in maltreated preschoolers. *Journal of Clinical Child Psychology*, 29, 307–318.

- Toth, S. L., Maughan, A., Manly, J. T., Spagnola, M., & Cicchetti, D. (2002). The relative efficacy of two interventions in altering maltreated preschool children's representational models: Implications for attachment. *Development and Psychopathology, 14*, 877-908.
- Toth, S. L., Rogosch, F. A., Manly, J. T., & Cicchetti, D. (2006). The efficacy of toddler-parent psychotherapy to reorganize attachment in the young offspring of mothers with major depressive disorder: A randomized preventive trial. *Journal of Consulting and Clinical Psychology, 74*, 1006-1016.
- Toth, S. L., Rogosch, F. A., Sturge-Apple, M., & Cicchetti, D. (2009). Maternal depression, children's attachment security, and representational development: An organizational perspective. *Child Development, 80*, 192-208.
- vonKlitzing, K., Stadelmann, S., & Perren, S. (2007). Story stem narratives of clinical and normal kindergarten children: Are content and performance associated with children's social competence? *American Psychologist, 9*, 271-286.
- Warren, S. L., Emde, R. N., & Sroufe, L. A. (2000). Internal representations: Predicting anxiety from children's play narratives. *Journal of the Academy of Child & Adolescent Psychiatry, 39*, 100-107.
- Warren, S. L., Mantz-Simmons, L., & Emde, R. N. (1993). *Narrative emotion coding*. Unpublished manuscript.
- Warren, S. L., Oppenheim, D., & Emde, R. N. (1996). Can emotions and themes in children's play predict behavior problems? *Journal of the American Academy of Child & Adolescent Psychiatry, 34*, 1331-1337.
- Weaver, T. L., & Clum, G. A. (1993). Early family environments and traumatic experiences associated with borderline personality disorder. *Journal of Consulting and Clinical Psychology, 61*, 1068-1075.
- Weiss, M., Zerkowitz, P., Feldman, R. B., Vogel, J., Heyman, M., & Paris, J. (1996). Psychopathology in offspring of mothers with borderline personality disorder. *Canadian Journal of Psychiatry, 41*, 285-290.
- Westen, D., & Cohen, R. P. (1993). The self in borderline personality disorder: A psychodynamic perspective. In Z. V. Segal & S. J. Blatt (Eds.), *The self in emotional distress; Cognitive and psychodynamic perspectives* (pp. 334-368). New York: Guilford Press.
- Winer, B. J., Brown, D. R., & Michels, K. M. (1991). *Statistical principles in experimental design* (3rd ed.). New York: McGraw-Hill.
- Yates, T. M. (2004). The developmental psychopathology of self-injurious behavior: Compensatory regulation in posttraumatic adaptation. *Clinical Psychology Review, 24*, 35-74.
- Yates, T. M., Carlson, E. A., & Egeland, B. (2008). A prospective study of child maltreatment and self-injurious behavior in a community sample. *Development and Psychopathology, 20*, 651-671.
- Zahn-Waxler, C., Schmitz, S., Fulker, D., Robinson, J. L., & Emde, R. N. (1996). Behavior problems in 5-year-old monozygotic and dizygotic twins: Genetic and environmental influences, patterns of regulation, and internalization of control. *Development and Psychopathology, 8*, 103-122.
- Zanarini, M. C. (2000). Childhood experiences associated with the development of borderline personality disorder. *Psychiatric Clinics of North America, 23*, 89-101.
- Zanarini, M. C., Frankenburg, F. R., Dubo, E. D., Sickel, A. E., Trikha, A., Levin, A., et al. (1998). Axis I comorbidity of borderline personality disorder. *American Journal of Psychiatry, 155*, 1733-1739.
- Zanarini, M. C., Frankenburg, F. R., Khera, G. S., & Bleichmar, J. (2001). Treatment histories of borderline inpatients. *Comprehensive Psychiatry, 42*, 144-150.
- Zanarini, M. C., Williams, A. A., Lewis, R. E., Reich, R. B., Vera, S. C., Marino, M. F., et al. (1997). Reported pathological childhood experiences associated with the development of borderline personality disorder. *American Journal of Psychiatry, 154*, 1101-1106.